

Chapter 17: Phenomena

Phenomena: Different masses of solute were added to 1 kg of either H_2O or C_6H_6 . The boiling and freezing points of the solutions were then measured. Examine the data to determine patterns in how these variables affect the boiling and freezing points of the solutions.

| H_2O | Exp. | Solute | Mass of Solute | Moles of Solute | Boiling Point | Freezing Point |
|----------|------|----------------------|----------------|-----------------|---------------|----------------|
| | 1 | None | 0.0 g | 0.00 mol | 100.0°C | 0.0°C |
| | 2 | NaCl | 50.0 g | 0.85 mol | 100.9°C | -3.2°C |
| | 3 | NaCl | 100.0 g | 1.71 mol | 101.7°C | -6.4°C |
| | 4 | K_2O | 50.0 g | 0.53 mol | 100.8°C | -3.0°C |
| | 5 | K_2O | 107.0 g | 1.14 mol | 101.7°C | -6.3°C |
| | 6 | $C_{12}H_{22}O_{11}$ | 1169.0 g | 3.42 mol | 101.7°C | -6.4°C |
| | 7 | $NaHCO_3$ | 71.7 g | 0.85 mol | 100.9°C | -3.2°C |
| | 8 | $C_6H_{12}O_6$ | 616.1 g | 3.42 mol | 101.7°C | -6.4°C |
| C_6H_6 | Exp. | Solute | Mass of Solute | Moles of Solute | Boiling Point | Freezing Point |
| | 1 | None | 0.0 g | 0.00 mol | 80.1°C | 5.5°C |
| | 2 | NaCl | 50.0 g | 0.85 mol | 84.4°C | -3.2°C |
| | 3 | NaCl | 100.0 g | 1.71 mol | 88.7°C | -12.0°C |
| | 4 | K_2O | 50.0 g | 0.53 mol | 84.1°C | -2.7°C |
| | 5 | K_2O | 107.0 g | 1.14 mol | 88.7°C | -11.9°C |
| | 6 | $C_{12}H_{22}O_{11}$ | 1169.0 g | 3.42 mol | 88.7°C | -12.0°C |
| | 7 | $NaHCO_3$ | 71.7 g | 0.85 mol | 84.4°C | -3.2°C |
| | 8 | $C_6H_{12}O_6$ | 616.1 g | 3.42 mol | 88.8°C | -12.0°C |

Chapter 17: Properties of Solutions

Chapter 17 Properties of Solutions

Big Idea: Liquids will mix together if both liquids are polar or both are nonpolar. The presence of a solute changes the physical properties of the system. For non-volatile solutes the vapor pressure, boiling point, freezing point, and osmotic pressure are only dependent on the number of ions/particles.

- Types of Solutions
- Solubility
- Colligative Properties
- Vapor Pressure
- Boiling and Freezing Point
- Osmotic Pressure

2

Types of Solutions

Matter can be broken in to two subcategories: pure substances and mixtures.

Pure Substances

- Elements:** A substance that cannot be separated into simpler components by chemical techniques.
- Compounds:** A specific combination of elements that can be separated into its elements by chemical techniques but not by physical techniques.

Examples: Water (H_2O), salt (NaCl), and sucrose ($C_{12}H_{22}O_{11}$)

3

Chapter 17: Properties of Solutions

Types of Solutions

Mixtures


- Heterogeneous Mixtures:** A mixture in which the individual components, although mixed together, lie in distinct regions that can be distinguished with an optical microscope.
- Homogeneous Mixture:** A mixture in which the individual components are uniformly mixed, even on the molecular scale.

Example: Air (nitrogen, oxygen, argon, ...)

4

Chapter 17: Properties of Solutions

Types of Solutions



5

Chapter 17: Properties of Solutions

Types of Solutions

Student Question

How many of the following are homogeneous mixtures?

- Oil and vinegar
- Chalk and table salt
- Salt water
- Kool-Aid
- Charcoal and sugar

- 1 is homogeneous
- 2 are homogeneous
- 3 are homogeneous
- 4 are homogeneous
- All are homogeneous

6

Chapter 17: Properties of Solutions

Types of Solutions

• **Solution:** A homogeneous mixture.

Solutions are made up of at least two parts

- **Solvent:** The most abundant component of a solution.
- **Solute:** A dissolved substance.

| State of Solution | State of Solvent | State of Solute | Example |
|-------------------|------------------|-----------------|--------------------------|
| Gas | Gas | Gas | Air Natural gas |
| Liquid | Liquid | Liquid | Tequila Antifreeze |
| Liquid | Liquid | Gas | Soda water |
| Liquid | Liquid | Solid | Sea water Sugar water |
| Solid | Solid | Solid | Alloys (Steel or brass) |
| Solid | Solid | Gas | Hydrogen in platinum |

Chapter 17: Properties of Solutions

Types of Solutions

Student Question


The density of a 40.0% by weight aqueous solution of NaOH is $1.432 \frac{g}{cm^3}$. What is the molality of NaOH?

Helpful Information: $M_{NaOH} = 40.00 \frac{g}{mol}$

- a) 12.9 m
- b) 14.3 m
- c) 16.7 m
- d) 13.8 m
- e) None of the above

Chapter 17: Properties of Solutions

Solubility



Chapter 17: Properties of Solutions

Solubility

Student Question


Which one of the following substances would be the most soluble in CCl_4 ?

- a) CH_3CH_2OH
- b) $C_{10}H_{22}$
- c) H_2O
- d) $NaCl$
- e) NH_3


Chapter 17: Properties of Solutions

Solubility

Cation and H_2O



Anion and H_2O



• **Hydration:** The reaction of a substance with water.


Note: For molecules, the extent of hydration increases as polarity increases.

Note: For ions, the extent of hydration increases as charge density (charge per volume) increases. In general, the smaller the size, the larger the charge density and the larger the charge, the larger the charge density.

Chapter 17: Properties of Solutions

Solubility


- **Saturated Solution:** A solution that holds the maximum amount of solute.
- **Unsaturated Solution:** A solution that holds less than the maximum amount of solute.
- **Supersaturated Solution:** A solution that holds more than the maximum amount of solute.



• If a solution is supersaturated any small disturbance can cause the solute to recrystallize

Chapter 17: Properties of Solutions

Solubility




- The graph shows the saturation level of different solutions at a given temperature. If a solution has a solubility that results in a point above the line, then the solution is considered to be supersaturated. However, if a solution has a solubility that results in a point under the line, then the solution is considered unsaturated.

13

Chapter 17: Properties of Solutions

Solubility



$$P = k_H \chi_{\text{gas}}$$

Note: Where χ is the mole fraction of the gas dissolved in solution.

Note: An alternate form of Henry's Law is: $P = k_H c$ where c is the molarity.

14

Chapter 17: Properties of Solutions

Colligative Properties

- Colligative Properties:** Physical properties of solutions that depend on the number of solute particles present but not the type of solute particles.
- Examples of Colligative Properties:**
 - Vapor Pressure (non volatile solutes)
 - Freezing Point
 - Boiling Point
 - Osmotic Pressure

15

Chapter 17: Properties of Solutions

Vapor Pressure

- Vapor Pressure:** The pressure exerted by the vapor of a liquid or solid.
- Volatile:** Having a high vapor pressure at ordinary temperatures (evaporates easily).
- Nonvolatile:** Having a low vapor pressure at ordinary temperatures (does not evaporate easily).

Nonvolatile Solute



- Raoult's Law**

$$P_{\text{solution}} = \chi_{\text{solvent}} P^{\circ}_{\text{solvent}}$$

Note: Where χ is the mole fraction of the solvent ($\frac{n_{\text{solvent}}}{n_{\text{total}}}$).

16

Chapter 17: Properties of Solutions

Vapor Pressure

Student Question

Which of the following aqueous solutions containing nonvolatile solutes should have the highest boiling point?


- 0.02 m $\text{C}_6\text{H}_{12}\text{O}_6$
- 0.02 m $(\text{NH}_4)_2\text{SO}_4$
- 0.02 m NaCl
- 0.02 m $\text{Ce}(\text{NO}_3)_4$
- All have the same boiling point

17

Chapter 17: Properties of Solutions

Vapor Pressure

Ideal Solution (Obeys Raoult's Law)



- Which substance (A or B) has the higher boiling point?

18

Chapter 17: Properties of Solutions

Vapor Pressure

Student Question

What is the vapor pressure of a solution of 50.0 g of CCl_4 and 50.0 g of CHCl_3 at 25°C . The vapor pressures at 25°C for pure CCl_4 and CHCl_3 are 98.3 torr and 172.0 torr respectively.

Helpful Information: $M_{\text{CHCl}_3} = 119.37 \frac{\text{g}}{\text{mol}}$ and $M_{\text{CCl}_4} = 153.81 \frac{\text{g}}{\text{mol}}$

- a) 131 torr
- b) 140. torr
- c) 149 torr
- d) 126 torr
- e) None of the Above


19

Chapter 17: Properties of Solutions

Vapor Pressure

Deviations from Raoult's Law

Ideal



$\Delta H=0$

20

Chapter 17: Properties of Solutions

Boiling and Freezing Points


- **Freezing Point Depression:** The decrease in the freezing point of a solvent caused by the presence of a solute.
- **Boiling Point Elevation:** The increase in the boiling point of a solution caused by the presence of a solute.

21

Chapter 17: Properties of Solutions

Boiling and Freezing Points

Boiling Point Elevation




- The presence of a nonvolatile solute lowers the vapor pressure of the solution, therefore, a higher temperature must be present in order for the vapor pressure of the solution to reach 1 atm (normal boiling point).

22

Chapter 17: Properties of Solutions

Boiling and Freezing Points

- At the freezing point, the solid and the liquid are in equilibrium.




23

Chapter 17: Properties of Solutions

Boiling and Freezing Points

Freezing Point Depression



24

Chapter 17: Properties of Solutions

Boiling and Freezing Points

Boiling and Freezing Point Data

| Solvent | Boiling Point (°C) | K_b ($^{\circ}\text{C}\cdot\text{kg}/\text{mol}$) | Freezing Point (°C) | K_f ($^{\circ}\text{C}\cdot\text{kg}/\text{mol}$) |
|-----------------------------------|--------------------|---|---------------------|---|
| H ₂ O | 100.0 | 0.51 | 0.0 | 1.86 |
| CCl ₄ | 76.5 | 5.03 | -22.99 | 30. |
| CHCl ₃ | 61.2 | 3.63 | -63.5 | 4.70 |
| C ₆ H ₆ | 80.1 | 2.53 | 5.5 | 5.12 |
| CS ₂ | 46.2 | 2.34 | -111.5 | 3.83 |
| C ₄ H ₁₀ O | 34.5 | 2.02 | -116.2 | 1.79 |
| C ₁₀ H ₁₆ O | 208.0 | 2.95 | 179.8 | 40. |

Chapter 17: Properties of Solutions

Boiling and Freezing Points

Student Question

A solution of 0.640 g of azulene in 100.0 g of benzene boils at 80.230°C. The boiling point of benzene is 80.100°C; the K_b is 2.53 $\frac{^{\circ}\text{C}\cdot\text{kg}}{\text{mol}}$. What is the molecular weight of azulene?


Helpful Information: The van't Hoff factor is 1 for azulene.

- 108 $\frac{\text{g}}{\text{mol}}$
- 117 $\frac{\text{g}}{\text{mol}}$
- 134 $\frac{\text{g}}{\text{mol}}$
- 99 $\frac{\text{g}}{\text{mol}}$
- None of the Above

Chapter 17: Properties of Solutions


Osmotic Pressure

- Osmosis:** The flow of solvent from a lower concentration solution through a semipermeable membrane to a higher concentration solution
- Osmotic Pressure:** The pressure needed to stop the flow of solvent through a semipermeable membrane.



Chapter 17: Properties of Solutions

Osmotic Pressure



Chapter 17: Properties of Solutions

Take Away From Chapter 17

- Big Idea:** Liquid will mix together if both liquids are polar or both are nonpolar. The presence of a solute changes the physical properties of the system. For non-volatile solutes the vapor pressure, boiling point, freezing points, and osmotic pressure are only dependent on the number of ions/particles.
- Types of Solutions**
 - Know the difference between homogeneous and heterogeneous mixtures
 - Homogeneous Mixtures: Uniformly mixed (air)
 - Heterogeneous Mixtures: Not uniformly mixed (chocolate chip cookie)

Numbers correspond to end of chapter questions.

Chapter 17: Properties of Solutions

Take Away From Chapter 17

- Types of Solutions (Continue)**
 - Know the definition of a solution
 - Solution: Homogeneous mixture
 - Be able to determine solution composition (15,16,17)
 - Molarity
 - $M = \frac{n_{\text{solute}}}{V_{\text{solution}}}$
 - Mole Fraction
 - $\chi_A = \frac{n_A}{n_{\text{total}}}$
 - Mass Percent
 - $\text{mass \% } A = \left(\frac{m_A}{m_{\text{total}}} \right) 100\%$
 - Molality
 - $m = \frac{n_{\text{solute}}}{m_{\text{solvent}}}$

Numbers correspond to end of chapter questions.

Chapter 17: Properties of Solutions

Take Away From Chapter 17

• Solubility

- Be able to predict whether two substances are soluble (33,34)
 - Like dissolves like
 - Two polar substances are soluble with each other
 - Two nonpolar substances are soluble with each other
 - A nonpolar and a polar substance are not soluble with each other
- Be able to predict the extent of hydration (32)
 - The more polar the ion/molecule the greater the hydration
 - The greater the charge density the greater the hydration (In general the smaller the ion/molecule the greater the hydration and the larger the charge the greater the hydration)
- Know the effects of temperature on solubility
 - In general the greater the temperature the higher the solubility of a solid in a liquid
 - The greater the temperature the lower the solubility of a gas in a liquid
- Know the effects of pressure on the solubility of a gas (39)
 - $P = k_H \chi$

Numbers correspond to end of chapter questions

Chapter 17: Properties of Solutions

31

Take Away From Chapter 17

• Colligative Properties

• Vapor Pressure

- Be able to calculate the vapor pressure of a solvent and a nonvolatile solute (42,44,47,79)
 - $P_{\text{solution}} = X_{\text{solvent}} P_{\text{solvent}}^*$
- Be able to calculate the vapor pressure of solvent and volatile solute (49,51)
 - $P_{\text{solution}} = X_{\text{solvent}} P_{\text{solvent}}^* + X_{\text{solute A}} P_{\text{solute A}}^* + \dots$
- Be able to construct mole fraction vs. vapor pressure plots (52,53,54,55,57)
 - Know what a positive deviation to Raoult's Law implies
 - $\Delta H > 0$: solute/solvent interaction are weaker than solute/solute interactions or solvent/solvent interactions
 - Know what a negative deviation of Raoult's Law implies
 - $\Delta H < 0$: solute/solvent interaction are stronger than solute/solute interactions or solvent/solvent interactions

Numbers correspond to end of chapter questions

Chapter 17: Properties of Solutions

32

Take Away From Chapter 17

• Boiling and Freezing Points

- Be able to determine the van't Hoff factor (i) and know how this factor affects colligative properties (13,81,82)
- Be able to calculate freezing point and boiling point of a solution (62,63,70,74,124)
 - $T_f = T_f^* - \Delta T_f$ or $T_b = T_b^* + \Delta T_b$
 - $\Delta T_f = i K_f m$ or $\Delta T_b = i K_b m$

• Osmotic Pressure

- Be able to calculate osmotic pressure (71,117)
 - $\Pi = i MRT$

Numbers correspond to end of chapter questions

Chapter 17: Properties of Solutions

33